

A Lyman Continuum Explorer - LyContEx

Stephan R. McCandliss

The Johns Hopkins University

Abstract.

GALEX has identified large numbers of extra galactic objects that are too faint to be observed by *FUSE*. These star forming systems are potential sources of Lyman continuum (Lyc) radiation and may contribute to the metagalactic background radiation that ionizes most of the baryons in the universe. By measuring the level of Lyc radiation leaking from these galaxies we can quantify the contribution of stars, relative to quasars, to the ionizing background at low redshift ($0.025 \lesssim z \lesssim 0.4$) and provide insight into the reionization processes at high redshift. A meter class instrument with a moderate spectral dispersion, wide-field, multi-object spectrograph can detect Lyc leak from a statistically significant number of these galaxies, if the average escape fraction $f_{esc} \gtrsim 1\%$. Such an instrument will have enormous discovery potential because of the large number of extra-galactic objects that exist within its grasp. LyContEx will be scientifically productive, synergistic with existing surveys, and can be developed into a far ultraviolet mission in the near future.

1. Introduction

For all its considerable success *FUSE* has three limitations. The spectral resolution is too low, the spectral resolution is too high and the étendue ($A_{eff}\Omega$, grasp, information content) is limited to a single object.

The challenge for future far ultraviolet missions is to identify compelling new science opportunities enabled by instrumentation that overcomes these limitations. Advances in high resolution spectroscopy ($R \geq 20000$) will require a very large telescope (Moos et al. 2004), a new approach to spectrograph design, and a multi-decadal timescale for development. A meter class, high étendue, moderate dispersion far ultraviolet spectrograph will have a shorter development timescale and an enormous potential for discovery because of the large number of extra galactic sources within its grasp too faint to be observed by *FUSE*. One new science opportunity enabled by such an instrument will be the search for the elusive Lyc radiation leaking from low redshift star-forming galaxies.

McCandliss et al. (2004) have detailed a new moderate dispersion, wide field, far ultraviolet spectroscopic survey instrument incorporating an array of micro-shutters (Moseley et al. 2000) designed to reach significantly lower sensitivities than *FUSE* and increase the information gathering power. Observations of objects with flux levels below the *FUSE* background equivalent flux (BEF $\sim 5 \times 10^{-15}$ ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$), face an exposure time “Wall” requiring increasingly deep exposures. A LyContEx mission with $R \sim 3000$ will yield a BEF $\sim 2 \times 10^{-17}$ ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$. Xu et al. (2004) finds > 850 gal (°) $^{-2}$ brighter than 23 mag $_{AB} = 3 \times 10^{-17}$ ergs cm $^{-2}$ s $^{-1}$ Å $^{-1}$ at 1540 Å from *GALEX* number counts (Figure 1). I show here a statistically significant number of these galaxies will have detectable levels of Lyc if the $f_{esc} \gtrsim 1\%$.




Figure 1. Differential number counts from *GALEX* observations. The cross hatched areas have been integrated to determine the total number of galaxies brighter than the limiting fluxes on the top of the figure. The integrated number of galaxies per $(^\circ)^2$ are printed under the cross-hatchings.

2. Why Seek Lyc Escaping from Galaxies?

The escape fraction is a key parameter widely used to “fine tune” the ionization history of the universe (Madau & Shull 1996). It is widely thought that the ionizing background that permeates metagalactic space is produced by quasars and star-forming galaxies (Madau & Shull 1996; Shull et al. 1999; Heckman et al. 2001), yet the contribution from star-forming galaxies has yet to be directly and convincingly detected. At low redshift upper limits to the escape fraction range from 5 - 50% (Leitherer et al. 1995; Hurwitz et al. 1997; Heckman et al. 2001; Deharveng et al. 2001). At high redshift Steidel et al. (2001) have claimed a tentative detection in $z = 3.4$ Lyman break galaxy composite spectrum.

A quantitative assessment of the contribution of galaxies to the ionizing background will require a large number of observations before a Lyc luminosity function can be established (Deharveng et al. 1997). Heckman et al. (2001) point out that determining the sources, strength and evolution of the metagalactic radiation field is crucial to understanding the fundamental properties of the IGM where the bulk of the baryons in the universe reside. Association of individual escape fractions within the mix of galactic morphological/spectral types found in the early universe, would allow the ionizing background at high redshift to be calculated semi-empirically at an epoch where starbursts are expected to dominate the reionization process. This would constrain the allowable fine tuning in ionization history models.

3. Predicting the Number of Detectable Lyc Emitters

GALEX FUV differential number counts of galaxies and Starburst99 models (Leitherer et al. 1999) of continuous star formation can be used to estimate the




Figure 2. A Starburst99 model (solar metallicity, Salpeter IMF, upper mass cutoff of $100 M_{\odot}$) is used to estimate the minimum observable *GALEX* FUV flux for a LyC flux at the background limit of the proposed LyContEx for average escape fractions 1 and 15%.

number of detectable galaxies emitting LyC photons. An escape fraction of 1 and 15% is imposed by applying the appropriate neutral hydrogen absorption model to a Starburst99 continuous star formation model. Offsetting the resulting spectra at F_{900} to the expected LyContEx BEF, the limiting *GALEX* FUV flux can be found. Integration of the differential number counts to these limiting F_{1540} fluxes provides the prediction of the total number of star-forming galaxies that should be detectable (see Figure 1). I find for the whole sky $\sim 10^5$ detectable galaxies if $f_{esc} = 1\%$ and $\sim 10^7$ galaxies if $f_{esc} = 15\%$. These numbers should be sufficient to allow the determination of a LyC luminosity function.

Acknowledgments. I wish to thank Erik Wilkinson, Karl Glazebrook, Ivan Baldry, Jeffrey Kruk and Warren Moos, Luciana Bianchi, Gerhardt Meurer, Bill Blair, Paul Feldman and Kevin France for useful discussions and their support. This work is supported through NASA grant NNG04WC03G to JHU.

References

- Deharveng, J.-M., et al. 2001, A&A, 375, 805
- Deharveng, J.-M., Faiesse, S., Milliard, B., & Le Brun, V. 1997, A&A, 325, 1259
- Heckman, T. M., et al. 2001, ApJ, 558, 56
- Hurwitz, M., Jelinsky, P., & Dixon, W. V. D. 1997, ApJ, 481, L31+
- Leitherer, C., et al. 1995, ApJ, 454, L19+
- Leitherer, C., et al. 1999, ApJS, 123, 3
- Madau, P., & Shull, J. M. 1996, ApJ, 457, 551
- McCandliss, S. R., et al. 2004, Proc. SPIE, Vol 5488, p. 709-718, astro-ph/0407464
- Moos, H. W., McCandliss, S. R., & Kruk, J. W. 2004, Proc. SPIE, Vol. 5488, p. 1-12
- Moseley, S. H., et al. 2000, in Proc. SPIE Vol. 4178, p. 51-58
- Shull, J. M., et al. 1999, AJ, 118, 1450
- Steidel, C. C., Pettini, M., & Adelberger, K. L. 2001, ApJ, 546, 665
- Xu, C. K., et al. 2004, to appear in the *GALEX* special edition of ApJ